

This article and any supplementary material should be cited as follows:
 Spencer RJ, Drag LL, Walker SJ, Bieliauskas LA. Self-reported cognitive symptoms following mild traumatic brain injury are poorly associated with neuropsychological performance in OIF/OEF veterans. J Rehabil Res Dev. 2010;47(6):521-530.
 DOI:10.1682/JRRD.2009.11.0181

Appendix 2. Survey Given to Medical Professionals

Please fill out the following demographic information:

Department and discipline (MD, RN, etc.): _____

Number of years in clinical practice: _____ (following Medical school/Nursing school/Graduate school) Do you regularly refer patients for neuropsychological testing? Yes No
 Approximately how many patients do you see per week? Please use the correlation table below as a general guideline when making your estimates. There are no right or wrong answers, we are only interested in your opinions.

1. Please estimate the correlation between a patient's self-report of their memory and their actual scores on tests of memory. _____
2. Please estimate the correlation between a patient's self-report of their processing speed/organizational abilities and their actual scores on tests of processing speed/organizational ability. _____
3. Please estimate the correlation between a patient's self-report of their attention/concentration and their actual scores on tests of attention/concentration.

Correlation	Percent of one score accounted for by the other score
1.00	100% (Perfect correspondence between scores)
0.90	81%
0.80	64%
0.70	49%
0.60	36%
0.50	25%
0.40	16%
0.30	9%
0.20	4%
0.10	1%
0.00	0% (No relationship between the scores)