

Appendix.

Scoring system of residual-limb quality. Maximum Points = 100. *Source:* Chakrabarty BK. An audit of the quality of the stump and its relation to rehabilitation in lower limb amputees.

Prosthet Orthot Int. 1998;22(2):136–46. [[PMID: 9747998](#)].

Wound	Healed	+10	Tenderness	None	+10
	Unhealed	-5		Moderate	+5
	Infected	-10		Severe	-10
Edema	None	+10	Proximal Joint	None	+10
	Minimal	+5		Contracture	<20°
	Significant	-5	>20°		-20
Scar	Fully Mobile	+10	Bone End Sculpted (& covered)	Satisfactory	+10
	< ¼ Adherent	-5		Acceptable	+5
	> ¼ < ½ Adherent	-6		Unsatisfactory	-10
	Adherent	-10		Bone end exposed	-20
Skin	Sensate	+6	Dog Ears	None	+6
	Insensate	-6		Minimal	0
	Insufficient	-6		Significant	-6
Length of Residual Limb	Suitable*	+10	Redundant Tissue	None	+6
	Acceptable	+5		Minimal	+3
	Unsuitable	-10		Significant	-6
Shape	Conical/Cylindrical	+6	Additional Scars or Other Factors	No	+6
	Bulbous	-6		Yes	-6

*For transtibial residual limb: 14–16 cm distal to knee joint line or 30 cm above ground level, whichever is longer.